

Artemisinin Gel

Topical Formula



Item # 74950

Available in jars of 1 oz (28 grams) cream

The Possible Benefits of Artemisinin Gel

- Formulated to provide a convenient way to utilize artemisinin topically*
- Provides high quality, independently tested artemisinin

Description

Artemisinin Gel contains artemisinin, or Qinghaosu, the active constituent of the herb *Artemisia annua* (sweet wormwood). Artemisinin Gel contains artemisinin in a concentrated suspension, and includes a driving agent to facilitate absorption and effectiveness.* Although to date no studies have been done on the use of artemisinin formulations on skin surfaces, preliminary clinical observations suggest it may be of value.*

Oral artemisinin has been thoroughly investigated. Good quality *Artemisia annua* contains 0.3-0.5% artemisinin, so pure artemisinin provides hundreds of times more activity than the whole herb itself. Our artemisinin has had independent cell tests verifying its effectiveness, and we do independent potency assays on every batch, using HPLC. In addition, we have received excellent feedback from numerous practitioners who utilize our Artemisinin capsules in their practice.*

Ingredients: Artemisinin, Purified Water, Poloxamer-407, Potassium Sorbate, Isopropyl Palmitate, Lecithin, Sorbic Acid, Ethoxydiglycol, Azelaic Acid.

Suggested Use: Apply 1/8 teaspoon (approx. 0.5 g) on clean skin only.

Caution: For external use only. Carefully clean hands and surface of skin before application. In rare cases artemisinin may cause idiosyncratic liver dysfunction.

Available to healthcare practitioners only.

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.

References

- Ames JR, Ryan MD, et al. *J. Free Rad Biol. Med* 1985, 1:353-61.
- Beekman AC, Wierenga PK, Woerdenbag HJ, Van Uden W, Pras N, Konings AW, el-Feraly FS, Galal AM, Wikstrom HV. *Planta Med.* 1998 Oct;64(7):615-9.
- Beekman AC, Woerdenbag HJ, Van Uden W, Pras N, Konings AW, Wikstrom HV. *J Pharm Pharmacol.* 1997 Dec;49(12):1254-8.
- Bharel S, Gulati M, Abdin P, Srivastava S. *Fitoterapia Vol LXVII No 5*, 1996.
- Borstnik K, Paik I, Shapiro TA, Posner GH. *Int J Parasitol.* 2002 Dec;32(13):1661-7.
- China Cooperative Research Group. *J. Trad. Chin. Med* 2, 17, 1982.
- Efferth T, Dunstan H, Sauerbrey A, Miyachi H, Chitambar CR. *Oncol.* 2001, Apr;18(4):767-73.
- Efferth T, Olbrich A, Bauer R. *Biochem Pharmacol.* 2002 Aug 15;64(4):617-23.
- Fishwick J, Edwards G, Ward SA, McLean WG. *Neurotoxicology.* 1998 Jun;19(3):393-403.
- Gulati A, Bharel S, Srivastava M, Abdin MZ. *Fitoterapia Vol LXVII No 5*, 1996.
- Keith Arnold, Tran Tinh Hien, et al. *Transactions of the Royal Society of Tropical Medicine and Hygiene* (1990) 84:499-502.
- Klayman D. *Science*, 1985, Vol. 238, May 31, p.1049
- Krungkrai SR, Yuthavong. *Trans. R Soc. Trop. Med Hyg* 1987, 81:710-4.
- Lee CH, Hong H, Shin J, Jung M, Shin I, Yoon J, Lee W. *Biochem Biophys Res Commun.* 2000 Aug 2;274(2):359-69.
- Levander OA, Ager AL, Morris VC. *Am.J. Clin. Nutr.* 1989; 5:346-52.
- Lu L. *Zhongguo Zhong Xi Yi Jie He Za Zhi.* 2002 Mar;22(3):169-71. Chinese.
- McLean WG, Ward SA. *Med Trop (Mars).* 1998;58(3 Suppl):28-31.
- Meshnick SR. *Int J Parasitol.* 2002 Dec;32(13):1655-60.
- Moore JC, Lai H, Li JR, Ren RL, McDougall JA, Singh NP, Chou CK. *Cancer Lett.* 1995 Nov 27;98(1):83-7.
- Mturi N, Musumba C, Wamola B, Ogutu B, Newton C. *CNS Drugs.* 2003;17(3):153-65.
- Mukanganyama S, Widersten M, Naik YS, Mannervik B, Hasler JA. *Int J Cancer.* 2002 Feb 10;97(5):700-5.
- Phan GT, de Vries PJ, Tran BQ, Le HQ, Nguyen NV, Nguyen TV, Heisterkamp SH, Kager PA. *Trop Med Int Health.* 2002 Oct;7(10):858-64.
- Posner GH, Northrop J, Paik IH, Borstnik K, Dolan P, Kensler TW, Xie S, Shapiro TA. *Bioorg Med Chem.* 2002 Jan;10(1):227-32.
- Posner GH, Paik IH, Sur S, McRiner AJ, Borstnik K, Xie S, Shapiro TA. *J Med Chem.* 2003 Mar 13;46(6):1060-1065.
- Posner GH, Ploypradith P, Parker MH, O'Dowd H, Woo SH, Northrop J, Krasavin M, Dolan P, Kensler TW, Xie S, Shapiro TA. *J Med Chem.* 1999 Oct 21;42(21):4275-80.
- Reungpatthanaphong P, Mankhetkorn S. *Biol Pharm Bull.* 2002 Dec;25(12):1555-61.
- Sidhu AB, Verdier-Pinard D, Fidock DA. *Science* 2002 Oct 4;298(5591):210-3.
- Singh NP, Lai H. *Life Sci.* 2001 Nov 21;70(1):49-56.
- Smith H, Crandall I, et al. *Mem Inst Oswaldo Cruz.* 1992;87 Suppl 3:303-12.
- Smith SL, Maggs JL, Edwards G, Ward SA, Park BK, McLean WG. *Neurotoxicology.* 1998 Aug-Oct;19(4-5):557-9.
- Stermitz FR, Scriven LN, Tegos G, Lewis K. *Planta Med.* 2002 Dec;68(12):1140-1.
- Sun WC, Han JX, Yang WY, Deng DA, Yue XF. *Zhongguo Yao Li Xue Bao.* 1992 Nov;13(6):541-3. Chinese.
- Walgate R. *Bull World Health Organ.* 2002;80(8):685-6.
- White NJ, Pongtavornpinyo W. *Proc R Soc Lond B Biol Sci.* 2003 Mar 7;270(1514):545-54.
- Wilairatana P, Krudsood S, Treeprasertsuk S, Chalermrut K, Looareesuwan S. *Arch Med Res.* 2002 Jul-Aug;33(4):416-21.
- Woerdenbag HJ, Moskal TA, Pras N, Malingre TM, el-Feraly FS, Kampinga HH, Konings AW. *J Nat Prod.* 1993 Jun;56(6):849-56.
- Wong JW, Yuen KH, Nagappan S, Shahul WS, et al. *J Pharm Pharmacol.* 2003 Feb;55(2):193-8.
- Zhai ZL, Xiao SH. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi.* 2001;19(3):182-5. Chinese.
- Zhang JT. *Therapie.* 2002 Mar-Apr;57(2):137-50.
- Zheng GQ. *Planta Med.* 1994 Feb;60(1):54-7.



Allergy Research Group®
2300 North Loop Road, Alameda, CA 94502
Phone: 800-545-9960 or 510-263-2000
Fax: 800-688-7426 or 510-263-2100
www.AllergyResearchGroup.com

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.