

DIM

Enhanced Delivery System



Item # 74140
120 Vegetarian Capsules

The Possible Benefits of DIM, a Dietary Supplement

- Promotes healthy estrogen metabolism and balance*
- Helps promote the conversion of estrogen to its beneficial, protective 2-hydroxyestrone metabolites and reduces production of genotoxic 16 α -hydroxyestrone*
- Stimulates detoxification enzyme systems*

Description

DIM (diindolylmethane) has been shown to help regulate and promote a more efficient metabolism of estrogen, and an optimal ratio of estrogen metabolites.*

DIM Enhanced Delivery System contains BioResponse DIM®, a unique formulation containing pure diindolylmethane, an indole. Indoles are plant compounds with health promoting properties, and are found in cruciferous vegetables such as broccoli, cabbage, cauliflower and Brussels sprouts.* The phytochemicals in cruciferous vegetables have been shown to beneficially affect the body's hormonal and detoxification systems, and epidemiological studies support the health benefits of consuming these vegetables.* DIM is a major active acid-catalyzed derivative of one of the phytochemicals in cruciferous vegetables, indole-3-carbinol (I3C). DIM is thought to be responsible for the health effects of dietary I3C.* DIM Enhanced Delivery System is a stable, bioavailable form of DIM, made possible through a proprietary delivery system. The formula is co-solubilized with phosphatidylcholine, and micro-

encapsulated in starch particles.

Research over the past thirty years has determined that healthy estrogen metabolism is closely linked to several healthy parameters in men and women, particularly some involving the breast, uterus, prostate and other reproductive tissue.* Genetics, excess weight, poor diet and other lifestyle factors may result in an imbalance of estrogen metabolites. Xenoestrogenic compounds, such as organochlorine pesticides, can also significantly disrupt healthy estrogen metabolism.

These estrogen disruptors alter estradiol hydroxylation metabolism producing a higher ratio of the genotoxic 16 α -hydroxyestrone (16 α -OHE1) to the safer and weaker estrogenic 2-hydroxyestrone (2-OHE1).* The genotoxic 16 α -OHE1 can potentially disrupt several normal cellular metabolic processes.* DIM promotes the conversion of estrogen to its beneficial, protective 2-hydroxyestrone metabolites and reduces production of genotoxic 16 α -OHE1.* Modulating these aspects of estrogen metabolism,

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.

particularly the production of 16 α -OHE1, may contribute to healthy aging.*

The mechanisms for DIM's health benefits primarily involve the induction of mixed function oxidases and phase II detoxification enzyme systems by the binding and activation of the arylhydrocarbon receptor (AhR).* Research using human breast cells (MCF-7) has shown that the binding of DIM® to the arylhydrocarbon receptor can result in rapid formation of the nuclear AhR complex and consequent induction of gene expression and

synthesis of cytochrome P450 detoxification enzyme (CyP450A1).* DIM® consequently produces increased levels of the protective hydroxylated estrogen 2-OHE1.* Some have suggested that DIM may also positively affect cellular signaling pathways.*

It is becoming increasingly apparent that DIM may provide an important mechanism for supporting successful aging despite the increasing levels of xenoestrogenic compounds in our modern world.*

Serving Size: 4 Capsules
Servings Per Container: 30

Amount Per Serving:

BioResponse DIM® (A patented diindolylmethane complex: starch, DIM 300 mg (25% min.), Vitamin E succinate, phosphatidylcholine (soy), silica)

Other ingredients: Hydroxypropyl methylcellulose, cellulose, L-leucine.

Suggested Use: As a dietary supplement, 2 to 4 capsules one or two times daily , or as directed by a healthcare practitioner.

References

- Bell MC, Crowley-Nowick P, Bradlow HL, et al. *Gynecol Oncol* 2000;78:123-9.
Ben-Jonathan N, Cooper RL, Foster P, et al. *Environ Health Perspect* 1999;107 Suppl 4:605-11.
Bradlow HL, Davis D, Sepkovic DW, et al. *Sci Total Environ* 1997;208:9-14.
Bradlow HL, Sepkovic DW, Telang NT, et al. *Ann N Y Acad Sci* 1999;889:204-13.
Chang YC, Riby J, Chang GH, et al. *Biochem Pharmacol* 1999;58:825-34.
Chen I, Safe S, Bjeldanes L. *Biochem Pharmacol* 1996;51:1069-76.
Davis DL, Telang NT, Osborne MP, et al. *Environ Health Perspect* 1997;105 Suppl 3:571-6.
Ge X, Fares FA, Yannai S. *Anticancer Res* 1999;19:3199-203.
Lake BG, Tredger JM, Renwick AB, et al. *Xenobiotica* 1998;28:803-11.
Michnovicz JJ, Adlercreutz H, Bradlow HL. *J Natl Cancer Inst* 1997;89:718-23.
Riby JE, Chang GH, Firestone GL, et al. *Biochem Pharmacol* 2000;60:167-77.
Sanderson JT, Slobbe L, Lansbergen GW, et al. *Toxicol Sci* 2001;61:40-8.
Shertzer HG, Senft AP. *Drug Metabol Drug Interact* 2000;17:159-88.
Wattenberg LW, Loub WD. *Cancer Res* 1978;38:1410-3.



Allergy Research Group®
2300 North Loop Road, Alameda, CA 94502
Phone: 800-545-9960 or 510-263-2000
Fax: 800-688-7426 or 510-263-2100
www.AllergyResearchGroup.com

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.