

PhytoArtemisinin

(Hypoallergenic)



Item # 74990
90 Vegetarian Capsules

"I have been very impressed by the efforts of ARG to perform both purity analysis and activity analysis on its artemisinin. For me, it was not even necessary, since I had already observed the awesome clinical efficacy of the product. I am grateful that this product is available and at a most reasonable price." - Robert Jay Rowen, M.D., Editor-in-Chief, Second Opinion

Description

PhytoArtemisinin contains pure artemisinin (Qinghaosu), the active constituent of the herb *Artemisia annua* (sweet wormwood) enhanced with phytosaponins. From the perspective of Traditional Chinese Medicine, artemisinin is cold.* The phytosaponins, which are warm, complement this characteristic of artemisinin, resulting in a more balanced formula than pure artemisinin alone.* The combination of artemisinin with phytosaponins may provide other synergistic benefits, including enhancing the activities of absorption and targeting.*

Gleditsia (*Gleditsia sinensis*), also called Chinese honey locust, is found in the woodlands of western China. Traditionally, all parts of the gleditsia plant have been used, and have variously been found to have anti-pruritic, astringent, expectorant, hemostatic,

purgative and other properties.* A 2002 study showed it may potentially reduce the release of histamine from mast cells.*

Pure artemisinin is the active constituent of the herb *Artemisia annua*. Good quality *A. annua* contains 0.3-0.5% artemisinin, so pure artemisinin provides hundreds of times more of the active constituent artemisinin than the whole herb. Research has shown artemisinin to be particularly beneficial in balancing the microbiology of the GI tract.*

Our artemisinin has had independent cell tests verifying its effectiveness, and we do independent HPLC potency assays on every batch. In addition, we have received excellent feedback from numerous practitioners who utilize artemisinin in their practice.*

Serving Size: 2 capsules
Servings Per Container: 45

Amount Per Serving:

Gleditsia (spine) extract (30% saponins)	700 mg
Artemisinin	100 mg

Other ingredients: Hydroxypropyl methylcellulose, cellulose, L-leucine.

Suggested Use: As a dietary supplement, 1 or 2 capsules one or two times daily before meals, or as directed by a healthcare practitioner. Sensitive individuals may want to take with food.

Caution: Not indicated for pregnant or nursing women. Long-term administration (greater than 1 month) should be monitored by a healthcare practitioner and include liver enzymes and hemoglobin testing. Combining with antioxidants or iron may theoretically decrease effectiveness. Detoxification reactions may be experienced by some individuals. In rare cases may cause idiosyncratic liver dysfunction.

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.

References

- Dai Y, Chan YP, Chu LM, Bu PP. *Biol Pharm Bull.* 2002 Sep;25(9):1179-82.
- Duke J A, Ayensu E S. 1985 ISBN 0-917256-20-4
Fogarty J E, Lutin M. Running Press ISBN 0-914294-92-X
- Shin TY, Kim DK. *Arch Pharm Res.* 2000 Aug;23(4):401-6.
- Stuart Rev G A. Taipei, Southern Materials Centre.
Yeung Him-Che. Institute of Chinese Medicine, Los Angeles 1985.
- Ames JR, Ryan MD, et al. *J. Free Rad Biol. Med* 1985, 1:353-61.
- Beekman AC, Wierenga PK, Woerdenbag HJ, Van Uden W, Pras N, Konings AW, el-Ferally FS, Galal AM, Wikstrom HV. *Planta Med.* 1998 Oct;64(7):615-9.
- Beekman AC, Woerdenbag HJ, Van Uden W, Pras N, Konings AW, Wikstrom HV. *J Pharm Pharmacol.* 1997 Dec;49(12):1254-8.
- Bharel S, Gulati M, Abdin P, Srivastava S. *Fitoterapia Vol LXVII No 5,* 1996.
- Borstnik K, Paik I, Shapiro TA, Posner GH. *Int J Parasitol.* 2002 Dec;32(13):1661-7.
- China Cooperative Research Group. *J. Trad. Chin. Med* 2, 17, 1982.
- Efferth T, Dunstan H, Sauerbrey A, Miyachi H, Chitambar CR. *Oncol.* 2001, Apr;18(4):767-73.
- Efferth T, Olbrich A, Bauer R. *Biochem Pharmacol.* 2002 Aug 15;64(4):617-23.
- Fishwick J, Edwards G, Ward SA, McLean WG. *Neurotoxicology.* 1998 Jun;19(3):393-403.
- Gulati A, Bharel S, Srivastava M, Abdin MZ. *Fitoterapia Vol LXV11 No 5,* 1996.
- Keith Arnold, Tran Tinh Hien, et al. *Transactions of the Royal Society of Tropical Medicine and Hygiene* (1990) 84:499-502.
- Klayman D. *Science*, 1985, Vol. 238, May 31, p.1049
- Krungkrai SR, Yuthavong. *Trans. R Soc. Trop. Med Hyg* 1987, 81:710-4.
- Lee CH, Hong H, Shin J, Jung M, Shin I, Yoon J, Lee W. *Biochem Biophys Res Commun.* 2000 Aug 2;274(2):359-69.
- Levander OA, Ager AL, Morris VC. *Am.J. Clin. Nutr.* 1989; 5:346-52.
- Lu L. *Zhongguo Zhong Xi Yi Jie He Za Zhi.* 2002 Mar;22(3):169-71. Chinese.
- McLean WG, Ward SA. *Med Trop (Mars).* 1998;58(3 Suppl):28-31.
- Meshnick SR. *Int J Parasitol.* 2002 Dec;32(13):1655-60.
- Moore JC, Lai H, Li JR, Ren RL, McDougall JA, Singh NP, Chou CK. *Cancer Lett.* 1995 Nov 27;98(1):83-7.
- Mturi N, Musumba C, Wamola B, Ogutu B, Newton C. *CNS Drugs.* 2003;17(3):153-65.
- Mukanganyama S, Widersten M, Naik YS, Mannervik B, Hasler JA. *Int J Cancer.* 2002 Feb 10;97(5):700-5.
- Phan GT, de Vries PJ, Tran BQ, Le HQ, Nguyen NV, Nguyen TV, Heisterkamp SH, Kager PA. *Trop Med Int Health.* 2002 Oct;7(10):858-64.
- Posner GH, Northrop J, Paik IH, Borstnik K, Dolan P, Kensler TW, Xie S, Shapiro TA. *Bioorg Med Chem.* 2002 Jan;10(1):227-32.
- Posner GH, Paik IH, Sur S, McRiner AJ, Borstnik K, Xie S, Shapiro TA. *J Med Chem.* 2003 Mar 13;46(6):1060-1065.
- Posner GH, Ploypradith P, Parker MH, O'Dowd H, Woo SH, Northrop J, Krasavin M, Dolan P, Kensler TW, Xie S, Shapiro TA. *J Med Chem.* 1999 Oct 21;42(21):4275-80.
- Reungpatthanaphong P, Mankhetkorn S. *Biol Pharm Bull.* 2002 Dec;25(12):1555-61.
- Sidhu AB, Verdier-Pinard D, Fidock DA. *Science* 2002 Oct 4;298(5591):210-3.
- Singh NP, Lai H. *Life Sci.* 2001 Nov 21;70(1):49-56.
- Smith H, Crandall I, et al. *Mem Inst Oswaldo Cruz.* 1992;87 Suppl 3:303-12.
- Smith SL, Maggs JL, Edwards G, Ward SA, Park BK, McLean WG. *Neurotoxicology.* 1998 Aug-Oct;19(4-5):557-9.
- Stermitz FR, Scriven LN, Tegos G, Lewis K. *Planta Med.* 2002 Dec;68(12):1140-1.
- Sun WC, Han JX, Yang WY, Deng DA, Yue XF. *Zhongguo Yao Li Xue Bao.* 1992 Nov;13(6):541-3. Chinese.
- Walgate R. *Bull World Health Organ.* 2002;80(8):685-6.
- White NJ, Pongtavornpinyo W. *Proc R Soc Lond B Biol Sci.* 2003 Mar 7;270(1514):545-54.
- Wilairatana P, Krudsood S, Treeprasertsuk S, Chalermrut K, Looareesuwan S. *Arch Med Res.* 2002 Jul-Aug;33(4):416-21.
- Woerdenbag HJ, Moskal TA, Pras N, Malingre TM, el-Ferally FS, Kampinga HH, Konings AW. *J Nat Prod.* 1993 Jun;56(6):849-56.
- Wong JW, Yuen KH, Nagappan S, Shahul WS, et al. *J Pharm Pharmacol.* 2003 Feb;55(2):193-8.
- Zhai ZL, Xiao SH. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi.* 2001;19(3):182-5. Chinese.
- Zhang JT. *Therapie.* 2002 Mar-Apr;57(2):137-50.
- Zheng GQ. *Planta Med.* 1994 Feb;60(1):54-7.



Allergy Research Group®
2300 North Loop Road, Alameda, CA 94502
Phone: 800-545-9960 or 510-263-2000
Fax: 800-688-7426 or 510-263-2100
www.AllergyResearchGroup.com

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.