

ThioDox[®]

Glutathione Complex

(Hypoallergenic)



Available in bottles of 90 tablets

Item # 72640

The Possible Benefits of ThioDox[®], a Dietary Supplement

- Supports liver detoxification, primarily phase II*
- Provides antioxidant protection*
- Supports immune function and production of glutathione*
- Facilitates the production of cellular energy*
- Enhances the effectiveness of other antioxidants*

Description

ThioDox[®] provides nutritional support for detoxification and immune function, and supplies nutrients shown to be especially supportive for liver and brain function.* L-glutathione (GSH), alpha-lipoic acid (ALA), and N-acetyl-L-cysteine (NAC) are three antioxidants, each powerful on its own; together they affect a host of critical cellular processes including oxidation and reduction reactions, detoxification reactions, and immune function.* Their actions are involved in glutathione production, scavenging of potentially harmful reactive free radicals, and primarily phase II liver detoxification.*

Glutathione (GSH) is a powerful tripeptide antioxidant that inhibits the formation of free radicals, and is thought to be the most important cellular antioxidant. It has been shown to help protect against damage from cigarette smoke, radiation, chemicals, heavy metals and X-rays, through its ability to protect the mitochondria and cellular nuclear structures.* GSH spares vitamin C, is involved in the regulation of DNA repair, and plays an important role in the detoxification systems of the body.* We always use L-glutathione in the preferred reduced form.

N-acetyl-L-cysteine (NAC) is the stable

form of the amino acid cysteine. It provides many of the same benefits as GSH, both through its own detoxifying and antioxidant function, and as a precursor of GSH.* NAC can help the liver to detoxify pollutants such as cigarette smoke, auto exhaust, agricultural chemicals, cyanide, and toxic metals such as mercury, arsenic, chromium and lead, and can potentially help control the toxicity of some chemotherapy treatments.* Studies suggest NAC may support respiratory function through its activity on viscous mucus.* NAC is involved with the regulation of homocysteine levels. Through their mitochondrial protective antioxidant activity, NAC and alpha-lipoic acid used in combination have been shown to support mental and physical performance in animal studies and in preliminary human studies.*

Alpha-lipoic acid (ALA, thioctic acid) is an important antioxidant that is both fat-soluble and water-soluble. Because of this, ALA is sometimes called the “universal” antioxidant. ALA provides antioxidant protection throughout the body. It facilitates the production of energy in cells and enhances the effectiveness of other antioxidants, regenerating vitamins C and E. It supports healthy function of the eyes, and blood glucose within normal levels.* ALA is a powerful support for the detoxification

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.

processes of the liver.* It can reduce the toxicity of mercury, cadmium, lead, and excess copper, manganese, and zinc.* ALA can also bring about increases in cellular glutathione levels.

Selenium, Tetrahydrofurfuryl Disulfide, Riboflavin-5-Phosphate, Vitamin C are included to support the overall effectiveness

of ThioDox[®]. Tetrahydro-furfuryl disulfide and riboflavin-5-phosphate are both sulfur enzymes, which can remain in their active, beneficial form (reduced) in the presence of glutathione. The enzyme glutathione reductase, which is involved with sulfur-reducing reactions, is selenium dependant. Vitamin C is provided as a natural preservative.

Serving Size: 1 Tablet

Servings Per Container: 90

Amount Per Serving:

Vitamin C (as Ascorbic Acid)	10 mg
Thiamine (as Thiamine Tetrahydrofurfuryl Disulfide)	5 mg
Riboflavin (as Riboflavin-5-Phosphate)	10 mg
Selenium (as L-Selenomethionine)	20 µg
N-Acetyl-L-Cysteine	250 mg
L-Glutathione (reduced)	200 mg
Lipoic Acid	50 mg

Other ingredients: Dicalcium phosphate, cellulose, magnesium stearate, silicon dioxide, stearic acid, ethyl vanillin.

Suggested Use: As a dietary supplement, 1 tablet one to four times daily with meals, or as directed by a healthcare practitioner.

Precaution: Lipoic acid can lower blood sugar levels. If you have diabetes or hypoglycemia, use this dietary supplement with caution.

Selected References

Ballatori N, et al. N-acetylcysteine as an antidote in methylmercury poisoning. *Environ Health Perspect.* May1998;106(5):267-71.
Berkson BM. Thioctic acid in treatment of hepatotoxic mushroom (Phalloides) poisoning. *N Engl J Med.* Feb1979;300(7):371.
de Blasio R, et al. N-acetylcysteine (NAC) in preventing nausea and vomiting induced by chemotherapy in patients suffering from inoperable non-small cell lung cancer (NSCLC). *Chest.* 1996;110(4,Suppl):103S.
Evans JL, Goldfine ID. Alpha-lipoic acid: a multifunctional antioxidant that improves insulin sensitivity in patients with type 2 diabetes. *Diabetes Technol Ther.* 2000;2(3):401-13.
Gurer H, Ozgunes H, Oztezcan S, Ercal N. Antioxidant role of alpha-lipoic acid in lead toxicity. *Free Radic Biol Med.* Jul1999;27(1-2):75-81.
Hagen TM, Liu J, Lykkesfeldt J, Wehr CM, Ingersoll RT, Vinarsky V, et al. Feeding acetyl-L-carnitine and lipoic acid to old rats significantly improves metabolic function while decreasing oxidative stress. *Proc Natl Acad Sci U S A.* Feb2002;99(4):1870-1875.
Liu J, Head E, Gharib AM, Yuan W, Ingersoll RT, et al. Memory loss in old rats is associated with brain mitochondrial decay and RNA/DNA oxidation: Partial reversal by feeding acetyl-L-carnitine and/or R-alpha-

lipoic acid. *Proc Natl Acad Sci U S A.* Feb2002;99(4):2356-2361.
Millman M, et al. Use of acetylcysteine in bronchial asthma--another look. *Ann Allergy.* Apr1985;54(4):294-6.
Packer L. Antioxidant Properties of Lipoic Acid and Its Therapeutic Effects in Prevention of Diabetes Complications and Cataracts. *Ann N Y Acad Sci.* Nov1994;738:257-64.
Plotzker R, Jensen DM, Payne JA. Case report. Amanita virosa acute hepatic necrosis: treatment with thioctic acid. *Am J Med Sci.* Mar1982;283(2):79-82.
Riise GC, et al. The intrabronchial microbial flora in chronic bronchitis patients: a target for N-acetylcysteine therapy? *Eur Respir J.* Jan1994;7(1):94-101.
Roes EM, Raijmakers MT, et al. Effects of oral N-acetylcysteine on plasma homocysteine and whole blood glutathione levels in healthy, non-pregnant women. *Clin Chem Lab Med* 2002 May;40(5):496-8.
Zappacosta B, Persichilli S, Mordente A, Minucci A, Lazzaro D, Meucci E, Giardina B. Inhibition of salivary enzymes by cigarette smoke and the protective role of glutathione. *Hum Exp Toxicol* 2002 Jan;21(1):7-11.



Allergy Research Group®
2300 North Loop Road, Alameda, CA 94502
Phone: 800-545-9960 or 510-263-2000
Fax: 800-688-7426 or 510-263-2100
www.AllergyResearchGroup.com

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.