

Olive Leaf Andrographis

Immunity and Circulation*

Olive Leaf Andrographis contains standardized extracts of olive tree leaves (*Olea europaea*) and andrographis (*Andrographis paniculata*). **Olive Leaf Andrographis** is enhanced with preformed vitamin A, and vitamin C.

Olive leaf possesses a variety of properties which support balanced intestinal microbiology and immunity.* Andrographis also has immune-supporting as well as adaptogenic properties.* Both botanicals also support the circulatory system, healthy cholesterol within normal levels, and normal blood sugar regulation.*



#72430
90 vegetarian tablets

Key Features

- Provides support for balanced aspects of the body's immune system*
- Supports healthy circulation and has multiple cardiogenic properties*
- May support aspects of blood sugar regulation within normal levels*
- Provides adaptogenic and antioxidant activities*



Olives and olive oil have long been a staple food in Mediterranean countries, and substances obtained from parts of the olive tree, including the leaves, have traditionally been used to support digestive and cardiovascular health, and for general well-being.* Olive oil is a key component of the Mediterranean diet, which we now know correlates with a decreased risk of coronary heart disease.

Our olive leaf extract is concentrated to minimum 18% oleuropein, the bitter principle of olives. Oleuropein supports aspects of the body's immune response, with strong microbial-balancing properties.* Modern research confirms this, and also shows that oleuropein supports the circulatory system, a normal inflammatory response, and has antioxidant, hepatoprotective, and neuroprotective properties.*

Research shows that olive leaf can promote the relaxation of vascular smooth muscles, protect low density lipoproteins from oxidation, and reduce production of the prostaglandin thromboxane A2.* Components found in olive leaf can potentiate glucose-induced insulin release and peripheral uptake of glucose.*

Andrographis (*Andrographis paniculata*) is a bitter herb commonly used in Asian medical traditions, and is known in the Ayurvedic system as the "King of Bitters".* It has immune-supporting and adaptogenic properties, as well as potential to support the circulatory system and normal blood sugar regulation.*

Andrographis has been shown to stimulate the nonspecific and antigen-specific immune response, while retaining the ability to reduce levels of cytokines and chemokines associated with immune system activation.* It has been beneficially studied in humans both acutely and long-term, and for situations of immune challenge, as well as autoimmunity, showing a remarkable balance in its effects.* In addition, andrographis has antioxidant properties involving Nrf2 (nuclear factor erythroid 2-related factor) activation.*

Vitamin A is necessary for normal functioning of the immune system, in particular the mucosal immunity of the cells that line the digestive and urinary tracts and the lungs.* These cells are a first line of defense.* Vitamin A is also important for proper development of lymphocytes.* Vitamin C provides support to a wide spectrum of body systems.*

Supplement Facts

Serving Size 1 Tablet
Servings Per Container 90

Amount Per Serving	% Daily Value*
Vitamin A (as 2000 IU of Retinyl Palmitate)	600 mcg RAE 66%
Vitamin C (as Ascorbyl Palmitate)	42 mg 47%
Olive (Leaf) Extract (min. 18% Oleuropein)	500 mg †
Andrographis (Aerial Part) Extract (50% Andrographolides)	120 mg †

† Daily Value not established.
* Percent Daily Value are based on a 2,000 calorie diet

Other ingredients: Microcrystalline cellulose, dicalcium phosphate, stearic acid, croscarmellose sodium, magnesium stearate, silicon dioxide.

Suggested Use: As a dietary supplement, 1 tablet one or two times daily with food, or as directed by a healthcare practitioner.

References:

Bennani-Kabchi N, et al. Therapie. Nov1999;54(6):717-23. Bisignano G, et al. J Pharm Pharmacol. Aug1999;51(8):971-4. Ficarra P, Ficarra R, de Pasquale A, Monforte MT, Calabro ML. Farmaco. 1991 Jun;46(6):803-15..

Gonzalez M, et al. Planta Medica. 1992;58:513-515. Hirschman SZ. Nat New Biol. Aug1972;238(87):277-9. Petroni A, et al. Thromb Res. Apr1995;78(2):151-60. Pieroni A, et al. Pharmazie. Oct1996;51(10):765-8.

Renis HE. Antimicrob Agents Chemother. Aug1975;8(2):194-9.

Ribeiro Rde A, Fiuzza de Melo MM, De Barros F, Gomes C,

Trolin G. J Ethnopharmacol. 1986 Mar;15(3):261-9.

Soret MG. Antimicrobial Agents and Chemotherapy. 1969;9:160-66.

Tassou CC, Nychas GJ, Board RG. Biotechnol Appl Biochem. 1991 Apr;13(2):231-7..

Visioli F, Vinceri FF, Galli C. Experientia. 1995 Jan 15;51(1):32-4.

Visoli F, et al. Life Sciences. 1994;55:1965-71.

Walter WM Jr, Fleming HP, Etchells JL. Appl Microbiol. 1973 Nov;26(5):773-6..

Zaruelo A, et al. Planta Med. Oct1991;57(5):417-9.

Kumar A, et al. Int J Res Pharm Chem. 2012;2(1):116-24.

Thakur AK, et al. J Tradit Complement Med. 2014 Dec 19;5(1):42-50.

Phunikhom K, et al. J Med Assoc Thai. 2015 Jul;98 Suppl 6:S41-7.

Zhang XF, Tan BK. Acta Pharmacol Sin. 2000 Dec;21(12):1157-64.

Puri A, et al. J Nat Prod. 1993 Jul;56(7):995-9.

Wong SY, et al. J Neuroinflam. 2016 Dec;13(1):34.

Li Y, et al. Evid Based Complement Alternat Med. 2017;2017:8248142.

Pan CW, et al. Oncotarget. 2017 Jun 20;8(25):41202-10.

Seo JY, et al. Mediators Inflamm. 2017;2017:5906189.

Barbaro B, et al. Int J Mol Sci. 2014 Oct; 15(10): 18508-18524.