

GI Flora

Contains four beneficial probiotic bacteria, *L. acidophilus*, *L. rhamnosus*, *L. casei* and *B. longum*, providing support for intestinal microbial balance in both the small and large intestines.*

An optimally functioning intestinal system is crucial to the health of the whole body. The human gastrointestinal tract harbors trillions of microorganisms, some beneficial to our health and some not. The cells that line the intestines, called villi, form a single layer that regulates digestion and absorbs the digested products. Friendly (probiotic) bacteria live attached to the villi, finding food and shelter, and in turn providing benefits to their host. Probiotic bacteria naturally occur in fermented foods, such as live culture yogurt and sauerkraut. Nobel Prize laureate Elie Metchnikoff observed in the 19th century that people in the Balkans who ate yogurt and other foods cultured with lactobacilli were longer-lived. He theorized that ingestion of lactobacilli could prolong life by competitively inhibiting nondesirable microbes.

Key Features

- Helps maintain a healthy intestinal probiotic balance*
- Supports the structure and functional integrity of the epithelial lining*
- May boost immune response and support resistance*
- Can produce vitamins, enzymes, and organic acids that support normal intestinal pH*



SKU #72461
90 vegetarian capsules

Trillions of probiotic microflora are found in the healthy small and large intestines, from up to 400 strains. They can support the structure and functional integrity of the epithelial lining by helping to metabolize vitamins, minerals and hormones, improve intestinal motility and assist in detoxification.* They can boost immune function, and have been shown to support resistance.* They produce metabolites such as lactic acid, hydrogen peroxide, bacteriocins and acetic acid that normalize the pH of the intestine and promote a healthy micro-ecological balance.* They support healthy conditions in the vagina, and cholesterol within normal levels.* They can produce lactase, the enzyme that digests lactose (milk sugar). When probiotics are depleted, supplemental probiotic bacteria are often needed in large amounts – in some cases, ten billion colony forming units (CFU) per day or more may be needed to promote intestinal balance.*

Members of the genus *Lactobacillus* take up residence primarily in the wall of the small intestine, where they provide many functions, including normalization of pH, promotion of digestive function, and stimulation of immune response.* *L. acidophilus* produces DL-lactic acid, amylase (carbohydrate-digesting enzyme), and bacteriocins. It is resistant to bile and gastric acids and has moderate lactase activity. *L. rhamnosus* has been shown to support the activity of both white blood cells and lymphocytic natural killer cells.* It also sometimes occurs in the large intestine, where it can create favorable conditions for the implantation of bifidobacteria. Compared to *L. acidophilus*, *L. rhamnosus* is less tolerant to bile and is more transient. It is a good releaser of lactase throughout the stomach and small intestine. *L. casei* can survive in a wide pH and temperature range, and it also produces DL-lactic acid and amylase. It is smaller than *L. acidophilus* and complements its growth. The name *casei* originates from the Latin word *caseus*, meaning cheese. All three of the above-mentioned lactobacilli were originally isolated from the human intestinal tract.

Almost 30 different species of *Bifidobacteria* have been identified, and they are the most plentiful probiotic bacterial group. They are more delicate than other common probiotics, and can easily be depleted by intestinal toxins or other stressors. Bifidobacteria are found in the large intestine and to a lesser extent in the lower part of the small intestine. *B. longum* was first isolated from human adults, and is strongly related to *B. infantis*, another well known probiotic. In breastfed infants bifido-bacteria comprise more than 95% of intestinal bacteria. They are anaerobic, and unlike other probiotic bacteria, they can ferment carbohydrates to both acetic and formic acids. They also produce lactic acid, creating a healthy pH in the colon.* They produce vitamins B1, B6, folic acid, and enzymes such as casein phosphatase and lysozyme. They also support the absorption of vitamins, and can take up iron from the contents of the bowel. They stimulate IgA production in the intestinal lining, bolstering the epithelial immune response.* They may help decompose potentially nonbeneficial metabolites.* Bifidobacteria are resistant to bile and gastric acids and have moderate lactase activity. The small amount of fructooligosaccharides (FOS) in GI Flora, naturally occurring long-chain carbohydrates that are not digested or absorbed by humans, can serve as a food for bifidobacteria.

Supplement Facts

Serving Size 3 Capsules
Servings Per Container 30

Amount Per Serving	% Daily Value
<i>Lactobacillus rhamnosus</i> UAL-r06™ (4.8 Billion CFU)	24 mg †
<i>Lactobacillus casei</i> UALc-03™ (4.8 Billion CFU)	16 mg †
<i>Bifidobacterium longum</i> UABl-14™ (1.2 Billion CFU)	12 mg †
<i>Lactobacillus acidophilus</i> La-14 (1.2 Billion CFU)	6 mg †

† Daily Value not established.

Other ingredients: Hydroxypropyl methylcellulose, FOS, rice flour, stearic acid.

Suggested Use: As a dietary supplement, 1 to 3 capsules two or three times daily, or as directed by a healthcare practitioner.

Stable at room temperature. Keep in a cool, dry place, tightly capped.